


Occupation and Mortality:
The National Health Interview Survey 1986-1994

Monograph



Lora E Fleming MD PhD, Terry Pitman BA, William LeBlanc PhD, Alberto Caban Jr MPH,
Orlando Gómez-Marín PhD MSc, David Lee PhD,
Diane Zheng MS, Dulce Jane MA

.....**Study Website: [http://www.I A\]Ua \]CF; 'Wca](http://www.IA]Ua]CF; 'Wca)**

Department of Epidemiology and Public Health
University of Miami School of Medicine
Miami, FL
33136

April 2005



Table of Contents

Abstract.....	3
Keywords	4
Acknowledgements	5
Introduction	6
Background.....	8
Methods	11
Result.....	13
Conclusions	18
References.....	21
Appendix.....	28

ABSTRACT

The National Health Interview Survey (NHIS) is a multipurpose household survey of the US civilian non-institutionalized population conducted annually since 1957. From 1986-1994, over 450,000 US workers, age 18 years and older, participated in a probability sampling of the entire non-institutionalized US population; variables collected included a range of measures of acute and chronic disability. Recently, the NHIS conducted a Mortality Follow-up with cause of death through 1997 available to researchers.

The objective of this Monograph was to review the mortality experience for US workers by occupation using the 1986-1994 NHIS data with follow-up through 1997. After adjustment for sample weights and design effects using SUDAAN, the unadjusted and age adjusted mortality rates (with their standard errors) using the direct method and the 2000 US Standard population were created in tabular format. The top 10 US causes of death identified by the Centers for Disease Control and Prevention (CDC) for the 1986-1994 Study Period were selected. These mortality rates have been presented by occupational subgroups, as well as by gender, race, and ethnicity subgroups within each occupation. Additional information has been made available to allow for extrapolation to the entire US worker population during the 1986-1994 time period. The Study Website (http://www.nchs.gov/nhis/occupational_mortality/) contains a repository of interactive tables which are available in both Excel and PDF formats; additional study and NHIS documentation are also available at this Website.

Understanding the occupational risk factors and improving the health of US workforce remains a public health priority. Occupational surveillance using the NHIS dataset allows for the careful monitoring of these occupational risk factors in order to identify prevention strategies and to minimize occupational mortality.

KEY WORDS

Mortality, Top 10 Causes of Death, Occupation, Industry, National Health Interview Survey (NHIS), Surveillance

ACKNOWLEDGMENTS

The data utilized in this publication were made available in part by the Inter-University Consortium for Political and Social Research. The data for the National Health Interview Survey (NHIS) were originally collected and prepared by the US Dept of Health and Human Services and the National Center for Health Statistics. Neither the collector of the original data nor the Consortium bears any responsibility for the analyses or interpretations presented in this publication. This study was funded in part through the National Institute of Occupational Safety and Health (NIOSH) Grant number 1 R01 OH03915-01. Additional information on this study can be found at the Study Website located at: <http://www.IAJUaJCF;Wca>.

INTRODUCTION

It is recognized that a variety of occupational and environmental risk factors interact to determine the overall health and well being of the US workforce. Occupational health surveillance is poised to analyze these interactions by systematically collecting, analyzing, and interpreting health data essential to the planning, implementation and evaluation of public health strategies to maximize workforce health. These robust surveillance systems collect and maintain health outcome information for all members of a temporally and geographically defined population at risk. The NIOSH NHIS Research Group at the University of Miami is funded by the National Institute for Occupational Safety and Health (NIOSH) to study the National Health Interview Survey (NHIS) dataset collected and conducted by the National Center for Health Statistics (NCHS) at the Centers for Disease Control (CDC).

The National Health Interview Survey (NHIS) is a continuous multipurpose and multistage probability area survey of the US civilian non-institutionalized population living at addressed dwellings (Kaminski 1980, Botman 1995, NCHS 2001). Each week a probability sample of households is interviewed by trained personnel to obtain information about the characteristics of each member of the household (Liao 1998). Data from the 1986-1994 NHIS Surveys included a range of measures of acute and chronic disability collected for all participants. Recently, the NHIS conducted a Mortality Follow-up with cause of death through 1997 for approximately 97% of the NHIS survey population from 1986-1994. The NHIS database allows for longitudinal analysis of mortality data as a retrospective cohort study, as well as for cross-sectional and trend analysis of the aggregate morbidity data. Thus, the NHIS database represents a unique opportunity to explore new research hypotheses, and to use the data as a surveillance tool to evaluate time trends, occupational disease, and mortality in the US for the past 2 decades in both genders and in a variety of race-ethnic subpopulations.

This Mortality Monograph establishes and applies a methodology to assess mortality rates by the top 10 causes of death for the US population during the 1986-1994 time period for US workers by occupation using the 1986-1994 NHIS data with follow-up through 1997. After adjustment for sample weights and design effects, the mortality rates were created in tabular format. These age adjusted mortality rates have been presented by occupation, as well as by gender, race, and ethnicity. Additional information has been made available to allow for extrapolation to the entire US worker population during the 1986-1994 time period.

BACKGROUND

The European countries, particularly England since 1837 in their Registrar General's Decennial Supplements for England and Wales, have had a long and illustrious history of performing nationwide occupational studies (Boffetta 1999, Drever 1995). As noted in the 1995 Registrar General's Report (Drever 1995), these data have provided a valuable means of generating hypotheses about work-related risks to health as well as insight in the effectiveness of preventive measures. The United States has had relatively few studies of equal scope and caliber to evaluate the causes of morbidity and mortality, and their trends, in US workers (Kaminski 1980, Milham 1976, Milham 1983, NIOSH 1993, Guralnick 1962, Guralnick 1963a, Guralnick 1963b, NIOSH 1997, Drever 1995, Gallagher 1989). As described below, the majority of these studies have focused on special subsets of data, not truly representative national data, and they have focused on mortality rather than morbidity.

US National Occupational Mortality Studies

Similar to the English studies, Guralnick (1962, 1963a, 1963b) used all US death certificates to look at age-race standardized mortality ratios (SMRs) for white and black men aged 20-64 by industry and by occupation. She evaluated trends by age, and compared them to the US Census data. Dr. Guralnick also acknowledged several study limitations, including what would now be called the healthy worker effect, and lack of information on important confounders such as socio-economic class. No female or other race-ethnic worker subpopulations were examined. Another major limitation was the use of death certificate-derived occupation/industry status; this has been shown to be highly inaccurate when compared to lifetime occupational histories, with misclassification estimated between 30-50% in US studies (Monson 1990, Checkoway 2004, Guralnick 1962, Guralnick 1963a, Guralnick 1963b, Kircher 1985, Schumacher 1986, Swanson 1984, Schade 1988).

Milham (1976, 1983) used data from Washington State to evaluate causes of mortality by occupation and industry using proportionate mortality ratios (PMRs) based on a large number of death certificates from 1950-1979. Limitations of these studies included no available confounders, the lack of rates (i.e. only proportions experienced), the lack of other race-ethnic worker subpopulations, lack of information on socio-economic class, and the use of death certificate information to classify occupation/industry status.

The National Traumatic Occupational Fatalities (NTOF) surveillance system estimates the risk of work-related fatal injuries for 50 industries and 50 occupations having the highest risks. NTOF supplies cause-specific risk estimates; it does not provide estimates for causes of death other than injury (Fosbroke 1997, Loomis 1998, Stout 1996). Similar analyses have been made using insurance databases (Toscano 1996). The National Institute of Occupational Safety and Health (NIOSH) (NIOSH 1993) used the death certificates with injury cause of death to evaluate fatal occupational injury rates from 1980-89 among male and female race-ethnic US worker populations. In addition to cause specific information, occupational injury rates (blacks>whites; males>females) and years of productive life lost were evaluated. Limitations to this study included no available confounders (such as socio-economic class), and the use of death certificate occupation/industry status.

Wagener et al (1997) attempted to use data from many available national databases (i.e., NHIS, NHANES, National Occupational Mortality Surveillance System, and NTOF, as well as the National Maternal and Infant Health Survey, National Hospital Ambulatory Medical Care Survey, the Census of Fatal Occupational Injuries, the Annual Survey of Occupational Injuries, and the Current Population Survey) to examine the health of women according to workforce and job conditions. In addition to presenting an overview of numbers and frequencies by age, race, ethnicity and educational level of various occupations, industries and presumed occupational exposures, for morbidity and mortality

analyses, the risks were calculated as the observed over expected proportions, not rates or trends over time.

Reviere et al (1995) used the National Mortality Follow-up Survey of 1986 to identify Sentinel Health Events Occupational (SHE(O)s) to look at causes of death beyond injury for the US worker. The National Cancer Institute (NCI), NIOSH and others have developed a database known as the National Occupational Mortality Surveillance System based on the death certificate listed occupation to study cancer and other causes of mortality in a variety of occupations (Ma 1998, Alterman 1997). NIOSH (1993, 1997) also evaluated mortality by occupation, industry, state and cause of death as proportionate mortality ratios from 24 reporting States from 1984-1988 among male and female black and white US worker populations. In addition to trends, years of productive life lost were estimated. The results may be evaluated by researchers and used as leads for further studies and to confirm previously identified associations. Researchers may identify new occupations and industries not previously recognized as experiencing an excess risk for a known occupational disease. Finally, these databases may be used to prioritize health promotion and intervention activities to the appropriate workers for both occupational and non occupational diseases (Kaminski 1980, Drever 1995). Limitations to these studies included no available confounders, the lack of rates and the use of only selected states' data, the lack of other race-ethnic worker subpopulations, and the use of death certificate occupation/industry.

Other Large Mortality and Morbidity Occupational Studies

In the US, as in other industrialized countries (Drever 1995), there have been a large number of studies of the mortality and morbidity rates of particular industries or groups of workers. Traditionally, occupational epidemiologic studies of mortality and morbidity have focused on cohorts of workers at particular industrial worksites, due to the relative ease of access to data (Honda 1995, Dell 1995, Tsai 1996, Chiazze 1997, Gold 1998). Research has also been performed on the causes of mortality and morbidity for white male workers in large occupational groups, such as chemical workers, construction workers, etc (Suruda 1996, Robinson 1995, Koskala 1997, Matanoski 1998, Steenland 1995, Blair 1993, Kross 1996, Savitz 1998). In addition, work has been done on morbidity and mortality trends for US workers in specific geographic areas (Fleming 1999, Hwang 1995, Milham 1976). Although performed in other countries (Kagamimori 1998, Engholm 1995, Schouten 1995), very little research has been performed on the US workforce as a whole, due to the lack of large and appropriately selected samples. Therefore, when studying the health status of the US worker, particular industry/worker group, and/or geographic constraints have applied.

Furthermore, there has been relatively little research on different age, race-ethnic and female subpopulations, as well as socio-economic class, of the US workforce as a whole. Recent Health disparity studies have shown that older age, race-ethnic, lower socio-economic class, and even some female subpopulations are at increased risk for disability and mortality compared to their white male counterparts (Pollan 1999, Zwering 1996 & 1998, King 1997, Bollini 1995, Murray 2003, Boffetta 1997, Barbeau 2004, Steenland 2003b, Steenland 2003a, Zwering 1997, O'Campo 2004, Janzen 2003, Khlal 2000, Frumkin 1999, Wagener 1991).

Although laudable, all of these attempts have been biased by selective reporting or by the use of occupation at time of death as the definition of occupational exposure, or have focused purely on traumatic injury, or are not generalizable to the entire US workforce due to sampling issues. As noted by Kaminski and Spirtas (1980), as a nationally representative dataset, the NHIS data can be used as a surveillance system for occupational disease morbidity and mortality for all US workers, and recommended that its use for this purpose be explored further.

METHODS

The NHIS dataset is anonymous and publicly available through the National Center for Health Statistics (NCHS); although there are no human subject considerations in the use of this dataset per the NCHS, an official waiver was obtained from the University of Miami School of Medicine Human Subjects Committee. During the 1986-1994 study period, annual NHIS survey response rates experienced ranged from 95-98% (Massey 1989). In the majority of cases (63%), the participants themselves answered all the questions, and for the remaining participants, the responses were obtained from their relatives or other proxies. For simplicity, in the present study either self-reported or proxy-reported data are referred to as “reported.”

Information on employment during the two weeks prior to the interview was collected for all persons 18 years or older in order to determine the person's employment status; as utilized by other investigators, all subjects age 18 years and older who had worked or experienced having jobs, both paid and unpaid, during the two weeks prior to the NHIS survey were considered currently employed (Kaminski 1980, Zwerling 1997, Brackbill 1988). Standardized Occupational Codes (really US Census codes) were provided in the NHIS database (NCHS 1998a, NCHS 1992) as well as various NHIS recodes with less detailed grouped occupational codes. For the purpose of this study, only the 206 occupations employing at least 100,000 workers annually in the US during the 1986-94 study period were selected for detailed analyses (including their US Census standardized occupational codes).

A Personal Identifier Code has been created by NCHS, allowing linkage with the individual NHIS survey years. The linkage was performed completely by computer, and used an algorithm with a match class and score. On average, the linkage was 97% complete; only those under 18 years and those with insufficient information to permit linkage to the NDI were considered ineligible and excluded from the analysis. With the NHIS Mortality Follow-up through December 31, 1997, a total of 12,124 deaths were found by this successful linkage with 440,245 working adults among the 451,897 who participated in the NHIS 1986-94 surveys (NCHS 1998b, NCHS 2001). This Mortality Follow-up permits the use of the NHIS Survey data since 1986 for survival, mortality, and life expectancy analyses. For example, Liao et al (1998) used a part of the NHIS Mortality Follow-up survey (1986-90) to examine mortality rates in Hispanics compared to Non Hispanic Whites in the US; the University of Miami Investigators have used the 1986-1994 Study Database to examine causes of mortality among all pesticide applicators and farmers compared to other US workers (Fleming 2003).

Provided in [Table 2](#) are the number of workers who participated in the NHIS survey pooled over the 1986-94 period (“sample size”), the number of NHIS worker participants with mortality follow-up through 1997 (“mortality follow-up sample size”), the estimated number of workers these participants represented in the US worker population during this time period (“estimated US worker population”), and the number of deaths are presented by gender, race and ethnicity subpopulations. Of note, tables for the 13, 41, and 206 occupational groups are available at the Study Website (www.UMiamiORG.com).

For the present analyses, the Investigators used the CDC Top 10 Causes of Death for the US population age 18 years and older from 1986-1994 ([Table 1](#)). The rankings of these causes of death were generated using the CDC program WISQARS TM (Web-based Injury Statistics Query and Reporting System) which is an interactive database system that provides customized reports of injury-related data. These data can be generated at the Website of the CDC National Center for Injury Prevention and Control: <http://webapp.cdc.gov/sasweb/ncipc/leadcaus9.html>. The ICD 9 Codes for these Top 10 Causes of Death can be found in [Table 1](#). For each of the Top 10 Causes of Death, analyses were performed by occupation and other factors available (i.e. gender, race, and ethnicity).

Statistical Methods

Because of the multi-stage sampling design, all analyses were performed with adjustment for sample weights and design effects using the SUDAAN statistical package (RTI 2001). The sample weights used were those required for the analysis of data from combined survey years and were calculated as specified by Botman and Jack (1995).

Age-adjusted mortality rates

Age-adjusted rates and the corresponding standard errors presented in the present monograph are expressed in terms of deaths per 100,000 person years, and were computed using the “direct method” using the year 2000 US population, ages 18 and older, as the standard population. The latter was obtained from the master list of age-adjustment weights, broken into 15 categories, from the CDC’s Healthy People 2010 Statistical Notes #20 (<http://www.cdc.gov/nchs/products/pubs/pubd/hp2k/statnt/20-11.htm>).

Using the Sudaan's “Proc Descript,” the investigator can perform calculation of age-standardized rates by the direct method with adjustments for sample weights and design effects. However, without suitable modification to the data set, this procedure does not allow the inclusion of person-years and does not take into account the changing age-structure of a cohort followed over time. Therefore, before using Sudaan's “Proc Descript,” the data set was rewritten so that there would be one record per year of follow-up for each individual, either until the year of death or until the end of the follow-up period, i.e., December 31, 1997. In addition, the survey respondent’s age on each additional record was advanced one year. These modifications generated a new data set that takes into account not only the number of years-at-risk for each survey participant, but also the age dynamics of the cohort.

Data Tables and Use

Mortality data are presented in tabular format for each of the 3 levels of major US occupational groupings (i.e. 13, 41, and 206). First, there is an initial table showing the age adjusted mortality rates (and their corresponding standard errors) per 100,000 for the study time period for overall mortality and the Top 10 Causes of Mortality for all workers of a particular occupational group ([Table 3](#)). Subsequent Tables present data on the age adjusted overall and cause specific mortality rates (and their corresponding standard errors) per 100,000 for the study time period for the different gender and race/ethnicity subpopulations with corresponding sample sizes.

Mortality rates were calculated when there was at least 1 death in the occupational subpopulation of interest. It should be noted that mortality analyses with less than 5 deaths/occupational subpopulation can result in very unstable point estimates and corresponding standard errors. Furthermore, because of the sample weighting of the NHIS, in some instances a very small standard error of approximately 0.0 will result. Therefore, analyses of small numbers of deaths should be interpreted with caution.

These unique data tables have been made available in two contemporary file formats: Portable Document Format (PDF) and Excel Spread Sheets (Excel) at the Study Website (URL: <http://www.UMiamiORG.com>). The PDF format allows researchers to quickly view and print the current table layouts, while the Excel format can be utilized to download the files to a remote computer and manipulate the data table locally.

To save an Excel file to a local computer system, the Study Website user can right click with their mouse over a link, at which time a dialog box will appear. Select the “Save target as” option to save the file from the Study website to the local system. A file viewer for PDF files is readily and freely available at the following website <http://www.adobe.com/products/acrobat/readstep2.html>.

Researchers can utilize these additional data tables to further explore mortality experienced among this population-based sample and extrapolate to the general US workforce. As discussed above, the Standard Errors (SEs) presented in the Tables for the age adjusted overall and cause specific mortality rates are presented. These SEs can be used to generate estimates of national level numbers as well as to generate confidence intervals. For example, for a particular disability measure, the reader can take $(1.96 \times SE) \pm \text{Mortality Rate}$ to generate the estimated range of that particular mortality rate among US workers.

RESULTS

The results presented below summarize the age adjusted overall and cause specific mortality rates (\pm their corresponding standard errors [SE]) per 100,000 persons enrolled during the study period 1986-1994 with follow-up through 1997 by describing the highest and lowest of the particular cause specific mortality rate for each of the occupational groupings (i.e. 13, 41, and 206). More detailed tables which include standard errors for the 13, 41 and 206 occupations are available at the Study Websites and Appendices as described above; in addition to occupation, all tables report the data by gender, race and ethnicity as well as total populations.

Overall Mortality

13 Occupations : In [Table 4](#), the age adjusted mortality rate for overall mortality by the different occupations is presented by gender, race and ethnicity, as well as the estimated US population numbers. Among the 13 occupations, Protective service workers experienced the highest overall mortality rates (1058 ± 150), while Technicians/related support workers experienced the lowest (454 ± 72). Among the gender-race-ethnicity subgroups, black Protective service workers experienced the highest overall mortality rate (2139 ± 678), while Hispanic Private household workers experienced the lowest (160 ± 72).

41 Occupations : In [Table 4](#), the age adjusted mortality rate for overall mortality by the different occupations is presented by gender, race and ethnicity, as well as the estimated US population numbers. Among the 41 occupations, Computer equipment operators experienced the highest overall mortality rate (1429 ± 869), while other transportation, except motor vehicles workers experienced the lowest (337 ± 103). Among the gender-race-ethnicity subgroups, Construction laborers in the 'other races' category experienced the highest overall mortality rate (4152 ± 2237), while eight subgroups of workers experienced no deaths.

206 Occupations : In [Table 4](#), the age adjusted mortality rate for overall mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Sales workers, parts (3111 ± 200) experienced the highest overall mortality, while Personnel and labor relations managers experienced the lowest (90 ± 32). Among the gender-race-ethnicity subgroups, black Industry workers experienced the highest overall mortality (6918 ± 803), while two hundred three worker subgroups experienced no deaths.

Cardiovascular Disease

In [Table 5](#), the age adjusted mortality rate for cardiovascular mortality by the different occupations is presented by gender, race and ethnicity, as well as the estimated US population numbers. Among the [13 occupations](#), Transportation/material moving workers experienced the highest rate of cardiovascular mortality (372 ± 93), while technicians/related support workers experienced the lowest (167 ± 52). Among the gender-race-ethnicity subgroups, black Executive, administrative, managerial workers experienced the highest cardiovascular mortality (714 ± 446), while Technicians/related support workers in the 'other races' category experienced no deaths.

41 Occupations : In [Table 5](#), the age adjusted mortality rate for cardiovascular disease by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Computer equipment operators experienced the highest rate of cardiovascular mortality (1251 ± 868), while Fabricators, assemblers, inspectors, samplers experienced the lowest (69 ± 15). Among the gender-race-ethnicity subgroups, Hispanic

Other professional specialty occupations experienced the highest rate of cardiovascular mortality (1313±864), while thirty subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 5 , the age adjusted mortality rate for cardiovascular mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Heating air conditioning, and refrigeration mechanics experienced the highest cardiovascular mortality rate (2155±38), while bank tellers and dental assistants experienced no deaths. Among the gender-race-ethnicity subgroups, Hispanic Medicine and health workers (4801±1615) experienced the highest cardiovascular mortality, while workers in four hundred ninety-five different subgroups of workers experienced no deaths.

Malignant Neoplasms

In [Table 6](#) , the age adjusted mortality rate for cancer mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the [13 occupations](#), Protective service workers experienced the highest cancer mortality rate (438±96), while technicians/related support experienced the lowest (139±25). Among the gender-race-ethnicity subgroups, black Protective service workers experienced the highest cancer mortality (1675±677), while private Household workers in the 'other races' category experienced no deaths.

[41 Occupations](#) : In Table 6 , the age adjusted mortality rate for cancer mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Forestry and fishing occupations experienced the highest rates of cancer mortality (755±237), while Computer equipment operators experienced the lowest (82±27). Among the gender-race-ethnicity subgroups, black Officials and administrators public administration experienced the highest cancer mortality (3961±113), while twenty-three subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 6 , the age adjusted mortality rate for cancer mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Sales workers experienced the highest rate of cancer mortality (2814±163), while operations/systems researchers and analysts experienced the lowest (17±17). Among the gender-race-ethnicity subgroups, Hispanic Garage and service station related occupations experienced the highest cancer mortality (5527±748), while four hundred eighteen subgroups of workers experienced no deaths.

Cerebrovascular Disease

In [Table 7](#) , the age adjusted mortality rate for cerebrovascular mortality by the different occupations is presented by gender, race, and ethnicity. Among the [13 occupations](#), Precision production, craft, repair workers experienced the highest rate of cerebrovascular mortality (78±28), while Technicians/related support experienced the lowest (4±2). Among the gender-race-ethnicity subgroups, black Handlers, equipment cleaners, helpers, laborers experienced the highest cerebrovascular mortality (169±97), while thirteen subgroups of workers experienced no deaths.

[41 Occupations](#) : In Table 7 , the age adjusted mortality rate for cerebrovascular mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Precision production occupations experienced the highest rates of cerebrovascular mortality (147±61), while Health technologists and technicians workers experienced the lowest (8±6). Among the gender-race-ethnicity subgroups, Mail and message distributing workers in the 'other races' category experienced the highest cerebrovascular mortality (375±266), while ninety-seven subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 7 , the age adjusted mortality rate for cerebrovascular mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Inspectors, testers, and graders experienced the highest cerebrovascular mortality rates (2473 ± 18), while seventy-four occupational groups experienced no deaths. Among the gender-race-ethnicity subgroups, Musicians and composers in the 'other races' category experienced the highest cerebrovascular mortality (6191 ± 0.0), while nine hundred forty-four subgroups of workers experienced no deaths.

Chronic Respiratory Disease

In [Table 8](#) , the age adjusted mortality rate for chronic respiratory disease mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the [13 occupations](#), Precision production, craft, repair workers experienced the highest chronic respiratory mortality rate (61 ± 25), while Private household workers experienced the lowest (5 ± 4). Among the gender-race-ethnicity subgroups, Hispanic Sales workers experienced the highest chronic respiratory disease mortality (75 ± 57), while twenty-three subgroups of workers experienced no deaths.

[41 Occupations](#) : In Table 8 , the age adjusted mortality rate for chronic respiratory disease mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Construction and extractive trades experienced the highest rate of chronic respiratory disease mortality (223 ± 122), while four occupational groups experienced no deaths. Among the gender-race-ethnicity subgroups, Construction laborers in the 'other races' category experienced the highest chronic respiratory disease mortality (484 ± 522), while one hundred twenty-one subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 8 , the age adjusted mortality rate for chronic respiratory disease mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Brickmasons and stonemasons experienced the highest rate of chronic respiratory disease mortality (546 ± 24), while ninety-eight occupational groups experienced no deaths. Among the gender-race-ethnicity subgroups, black Industrial workers experienced the highest chronic respiratory disease mortality (5517 ± 0.0), while one thousand seventy subgroups of workers experienced no deaths.

Unintentional Injury

[13 Occupations](#) : In [Table 9](#) , the age adjusted mortality rate for unintentional injury mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 13 occupations, transportation/ material moving workers experienced the highest rate of unintentional injury mortality (231 ± 82), while Executive, administrative managerial occupations experienced the lowest (21 ± 3). Among the gender-race-ethnicity subgroups, white Transportation/material moving workers experienced the highest unintentional injury mortality (342 ± 133), while seven subgroups of workers experienced no deaths.

[41 Occupations](#) : In Table 9 , the age adjusted mortality rate for unintentional injury mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Material moving equipment operators experienced the highest rate of unintentional injury mortality (563 ± 13), while Other transportation, except motor vehicles, workers experienced no deaths. Among the gender-race-ethnicity subgroups, male Material moving equipment operators experienced the highest unintentional injury mortality (565 ± 13), while seventy subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 9 , the age adjusted mortality rate for unintentional injury mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Real estate sales occupations experienced the highest rate of unintentional injury mortality (571±550), while forty-five occupational groups experienced no deaths. Among the gender-race-ethnicity subgroups, black Tool and die makers experienced the highest unintentional injury mortality (2696±2589), while seven hundred sixty-four subgroups of workers experienced no deaths.

Pneumonia and Influenza

In [Table 10](#) , the age adjusted mortality rate for pneumonia and influenza mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the [13 occupations](#), Machine operators, assemblers, inspectors experienced the highest rates of pneumonia and influenza mortality (65±56), while Technicians/related support workers experienced the lowest (1±1). Among the gender-race-ethnicity subgroups, male Machine operators, assemblers, inspectors experienced the highest pneumonia and influenza mortality (96±85), while nineteen subgroups of workers experienced no deaths.

[41 Occupations](#) : In Table 10 , the age adjusted mortality rate for pneumonia and influenza mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Machine operators/tenderers (except precision) experienced the highest rates of pneumonia and influenza mortality (79 ±67), while seven occupations experienced no deaths. Among the gender-race-ethnicity subgroups, male Machine operators/tenderers (except precision) experienced the highest pneumonia and influenza mortality (129±112), while one hundred and thirty-one subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 10 , the age adjusted mortality rate for pneumonia and influenza mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Roofers experienced the highest rates of pneumonia and influenza mortality (1227±0.0), while one hundred and fifteen occupations experienced no deaths. Among the gender-race-ethnicity three subgroups, white, male, and non-Hispanic Roofers experienced the highest pneumonia and influenza mortality (1227±0.0), while eleven hundred twenty-six subgroups of workers experienced no deaths.

Diabetes

In [Table 11](#) , the age adjusted mortality rate for diabetes mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the [13 occupations](#), Administrative support occupations (including clerical) experienced the highest rates of diabetes mortality (35±14), while Handlers, equipment cleaners, helpers, laborers experienced the lowest (4±2). Among the gender-race-ethnicity subgroups, Hispanic Protective service workers experienced the highest diabetes mortality (157±116), while twenty worker subgroups experienced no deaths.

[41 Occupations](#) : In Table 11 , the age adjusted mortality rate for diabetes mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Other Transportation (except motor vehicles) occupations experienced the highest rates of diabetes mortality (279±278), while seven occupations experienced no deaths. Among the gender-race-ethnicity subgroups, Hispanic Police and firefighters experienced the highest diabetes mortality (4436±2717), while one hundred twenty-five subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 11 , the age adjusted mortality rate for diabetes mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Supervisors, mechanics, and repairers experienced the highest rates of diabetes mortality (4211±1087), while one hundred and four occupations experienced no deaths. Among the gender-race-ethnicity subgroups, Hispanic Sheriffs/bailiffs/other law enforcement officers experienced the highest diabetes mortality (10755±0.0), while one thousand seventy-eight subgroups of workers experienced no deaths.

Suicide

In [Table 12](#) , the age adjusted mortality rate for suicide mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the [13 occupations](#), Handlers, equipment cleaners, helpers, laborers experienced the highest rate of suicide mortality (128±65), while private household workers experienced the lowest (18±18). Among the gender-race-ethnicity subgroups, Private household workers in the 'other races' category experienced the highest suicide mortality (639±685), while nineteen subgroups of workers experienced no deaths.

[41 Occupations](#) : In Table 12 , the age adjusted mortality rate for suicide mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Forestry and fishing occupations experienced the highest rate of suicide mortality (168±110), while Architects and surveyors experienced no deaths. Among the gender-race-ethnicity subgroups, Hispanic Personal service workers experienced the highest suicide mortality (826±763), while one hundred nine subgroups experienced no deaths.

[206 Occupations](#) : In Table 12 , the age adjusted mortality rate for suicide mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Legal assistants experienced the highest rates of suicide mortality (721±468), while eighty-two occupations experienced no deaths. Among the gender-race-ethnicity subgroups, Supervisors, cleaning and building service workers in the 'other races' category experienced the highest suicide mortality (2933±2650), while one thousand fifteen subgroups of workers experienced no deaths.

Liver Disease

In [Table 13](#) , the age adjusted mortality rate for liver disease mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the [13 occupations](#), Handlers, Equipment cleaners, helpers, laborers experienced the highest rate of liver disease mortality (123±42), while Administrative support occupations, including clerical workers experienced the lowest (14±6). Among the gender-race-ethnicity subgroups, Handlers, equipment cleaners, helpers, laborers in the 'other races' category experienced the highest liver disease mortality (244±240), while sixteen subgroups of workers experienced no deaths.

[41 Occupations](#) : In Table 13 , the age adjusted mortality rate for liver disease mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Other protective service occupations experienced the highest rate of liver disease mortality (152±87), while eight occupations experienced no deaths. Among the gender-race-ethnicity subgroups, Construction laborers in the 'other races' category experienced the highest liver disease mortality (4036±2931), while one hundred thirty-six subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 13 , the age adjusted mortality rate for liver disease mortality by the

different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Molding and casting machine operators experienced the highest rate of liver disease mortality (1036 ± 954), while one hundred sixteen occupations experienced no deaths. Among the gender-race-ethnicity subgroups, Construction laborers in the 'other races' category experienced the highest liver disease mortality (4036 ± 2931), while eleven hundred twenty-eight subgroups of workers experienced no deaths.

HIV/AIDS

In [Table 14](#) , the age adjusted mortality rate for HIV/AIDS mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the [13 occupations](#), Handlers, equipment cleaners, helpers, laborers experienced the highest rates of HIV/AIDS mortality (86 ± 25), while Farming, forestry, fishing workers experienced the lowest (24 ± 10). Among the gender-race-ethnicity subgroups, black Handlers, equipment cleaners, helpers, laborers experienced the highest HIV/AIDS mortality (330 ± 138), while thirteen subgroups of workers experienced no deaths.

[41 Occupations](#) : In Table 14 , the age adjusted mortality rate for HIV/AIDS mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 41 occupations, Architects and surveyors experienced the highest rates of HIV/AIDS mortality (226 ± 142), while two occupations experienced no deaths. Among the gender-race-ethnicity subgroups, black Construction and extractive trade workers experienced the highest HIV/AIDS mortality (599 ± 355), while eighty-six subgroups of workers experienced no deaths.

[206 Occupations](#) : In Table 14 , the age adjusted mortality rate for HIV/AIDS mortality by the different occupations is presented by gender, race and ethnicity as well as the estimated US population numbers. Among the 206 occupations, Designers experienced the highest rates of HIV/AIDS mortality (362 ± 192), while seventy-six occupations experienced no deaths. Among the gender-race-ethnicity subgroups, black Advertising and related sales occupations experienced the highest HIV/AIDS mortality (5991 ± 3917), while nine hundred forty-five subgroups of workers experienced no deaths.

CONCLUSIONS

The current Mortality Monograph of all currently employed adults 18 years or older from the 1986-1994 NHIS surveys with mortality follow-up for 97% through 1997 demonstrated increased overall and cause specific mortality rates among certain occupational groups compared to all others. This Monograph confirms some possible associations between particular occupations (and their exposures) and certain cause specific mortality rates already recognized in the literature. This Monograph also demonstrates possible new associations between occupations and cause specific mortality rates, particularly among the gender, race, and ethnic subpopulations with the more detailed occupational coding. In general, these new associations were only able to be detected due to the large representative sample size of the US workforce provided by the NHIS surveys. Furthermore, these possible new associations can be considered “hypothesis generating” and worthy of future investigation and research.

In general, among the occupations, more skilled “white collar” workers were less likely to have increased overall or cause specific mortality. Of note, these same workers were more likely to report less disability, better self rated health and fewer health conditions (Fleming 2004). Within particular occupations, Hispanic, black and “other” workers were more likely to experience increased overall and cause specific mortality. Of note, these same worker subgroups were more likely to report acute and chronic disability measures, poor self rated health and numbers of health conditions. In particular, race-specific comparisons often revealed higher morbidity in workers in the 'other races' category (Fleming 2004).

In a separate analysis, the investigators have modeled acute and chronic disability adjusting for age, gender, race/ethnicity, and education in the entire NHIS worker database, as well as comparing disability among farmers and pesticide applicators with all other US workers (Gomez Marin 2004, Fleming 2004). Among all NHIS workers, for acute disability younger age was a significant risk factor; older age was a significant risk factor in a dose response fashion for reporting chronic disability and poor health status. Of note, using the same NHIS database, Zwerling et al (1996, 1997, 1998) have found a significantly increased experienced injury risk among older and disabled US workers. As with other investigators, Gomez Marin et al (2004) also found that older female uneducated workers were at significantly higher risk of chronic disability and experienced poor health status compared to all other US workers. Less educated older female workers are more likely to have poorly paying jobs with significant physical labor and hours.

Several causes of mortality among those examined in this Monograph have been traditionally associated with occupational exposures: Malignant Neoplasms, Chronic Respiratory Disease, Pneumonia and Influenza, and Unintentional Injury (Nathell 2000, Rosengren 2004, Simpson 1999, Carpenter 1999, Rosenstock 2005). For Malignant Neoplasms, Forestry and fishing occupations had an increased mortality rate as has been explored in the literature with possible associations with exposure to certain herbicides and pesticides (Kirkhorn 2002, Milham 1999); it is interesting that the highest mortality rates were experienced by Hispanic Garage and service station related occupations. For Chronic Respiratory Disease, Construction and extractive trades had an increased rate also reported in the literature (Bernstein 2004, Bergdahl 2004), but the highest rates were seen among the black Industrial workers. For Pneumonia and influenza, Machine operators/tenderers (except precision) had elevated rates which is interesting given exposures to machining fluids with possible bacterial contamination and pulmonary effects (Anon 2002, Kreiss 1997), however the highest rates were seen among Roofers (particularly non Hispanic) who also experience pulmonary effects from occupational exposures and are members of the construction trades (Melius 2003). Finally for Unintentional Injury, as might be expected, Transportation/material moving workers had increased

mortality rates (Lam 2004, Janicak 2003), but the highest rates were seen among black Tool and die makers.

The other causes of mortality among the Top 10 Causes of Mortality in the US during the Study Period have had more tenuous associations with occupations and their exposures. For Overall Mortality, it is not surprising the Protective service workers (particularly black) and Construction workers (particularly “other races”) experienced increased mortality rates (Plani 2003, Dong 2004, Steenland 2003a, Steenland 2003b), but black Industry workers experienced the highest rates. With respect to Cardiovascular Disease, as might be expected, Transportation/material moving workers had increased rates (Martins 2003, Stein 2001), however Hispanic Medicine and health workers experienced the highest rates. For Cerebrovascular Disease, Precision production, craft, repair workers had increased mortality rates with the highest rates seen among Musicians and composers of “other races.” Diabetes Mortality rates were highest among Hispanic Sheriffs/bailiffs/other law enforcement officials. For Suicide Mortality, elevated mortality rates were seen among Forestry and fishing occupations possibly associated with the neurotoxicity of occupational pesticide exposures (Gregoire 2002, Parron 1996), however the highest rates were seen among Supervisors, cleaning and building service workers of “other races.” Liver Disease Mortality was highest among Construction workers of “other races.” Finally, HIV/AIDS Mortality was elevated among Designers (Cassano 2000), but highest among black Advertising and related sales occupations.

Limitations

These analyses suffered from some of the data limitations seen in similar previous epidemiologic studies. The most important limitation in interpreting the Mortality Rates was the relatively small number of deaths experienced to date by this NHIS Occupational Cohort, particularly those in the 206 Occupational Subgroups further divided by gender, race and ethnicity; therefore, as noted above, any increased rates must be viewed as hypothesis generating only. Other limitations included: the self report and cross sectional nature of the initial data which might lead to underestimation of the true health situation since really sick people leave the work force (part of the healthy worker effect); lack of individual exposure measures, and occupational misclassification in general due to the use of “current job” as a surrogate for “longest held job,” as well as misclassification of occupation related exposures. With regards to the latter issue, two occupational health related supplements were administered to NHIS participants, one in 1986 and the second in 1988. We used previously published Kappa values from these two supplements to estimate the extent to which “current job” reflected “longest held job” within each occupational category (Gomez Marin 2005). The use of the US worker population as the major comparison population is appropriate for controlling for the healthy worker effect and other biases (Cooper 1993, Checkoway 2004, Monson 1990, Burnett 1989). Previous work using the NHIS database has shown that certain occupational groups, such as farmers, smoke and drink less than many other worker groups (15-29% prevalence) which may explain some but not all of the study findings (Brackbill 1988, Nelson 1994, Sterling 1990, Lee 2004). Confounding data such as tobacco use and obesity are available for subpopulations of the NHIS database but were not used in this study; however, tobacco use data are available for use by researchers at the Study Website (www.UMiamiORG.com) (Caban in press, Lee 2004). The NHIS data (including occupation) were based largely on self-report (by the individual or proxy) without objective confirmation. The limitations and lack of these data are an important limitation on any conclusions that can be drawn from these analyses.

Although research in this area is often contradictory, validation studies conducted by the NCHS and others suggest that proxy reports lead to slightly lower prevalence estimates of a range of information compared with reports obtained directly from respondents (Edwards 1994, Edwards 1996, Thornberry 1987). To address this potential bias, we repeated our mortality analyses, including only the 63% of NHIS participants who were interviewed directly in the 1986-1994 NHIS surveys. For example,

findings indicated that the non-proxy self experienced responses of the disability measures were uniformly only slightly greater than those from the proxy responses (Fleming 2004).

Ultimately, the NHIS data strongly suggest that selected occupations by gender and race-ethnicity subpopulation should be an important target populations for occupational prevention and intervention (Partenen 2002). Researchers in occupational health are urged to seek out this important and publicly available resource. The NHIS database and mortality follow-up represent a probability sample of the entire US population, with the ability to compare both morbidity and mortality among US workers. Furthermore, as noted by NIOSH (1997) and the English Registrar's Decennial Reports (Drever 1995), databases such as the NHIS surveys and mortality follow-up can be used not only to target studies of work-related conditions and to add to the body of evidence generated from epidemiologic studies, but also to provide surveillance data for establishing priorities, and for tracking progress towards the elimination of preventable diseases and mortality.

REFERENCES

- Anonymous. From the Centers for Diseases Control and Prevention. Respiratory illness in workers exposed to metalworking fluid contaminated with nontuberculous mycobacteria-- Ohio, 2001. *JAMA*. 2002;287(23):3073-4.
- Alterman T, Burnett C, Peipins L, Lulich N, Halperin W. Occupation and cervical cancer: an opportunity for prevention. *J Womens Health* 1997;6:649-57.
- Barbeau E, Krieger N, Soobader M-J. Working class matters: socio-economic disadvantage, race/ethnicity, gender, and smoking in the NHIS 2000. *AM J Pub Health* 2004;94:269-278.
- Bergdahl IA, Toren K, Eriksson K, Hedlund U, Nilsson T, Flodin R, Jarvholm B. Increased mortality in COPD among construction workers exposed to inorganic dust. *European Respiratory Journal*. 2004;23(3):402-6.
- Bernstein DM. Increased mortality in COPD among construction workers exposed to inorganic dust. *European Respiratory Journal*. 2004;24(3):512.
- Blair A, Dosemeci M, Heineman E. Cancer and other causes of death among male and female farmers from 23 states. *Am J Ind Med* 1993;23:729-42.
- Boffetta P, Kogevinas M, ed. Occupational Cancer in Europe. *Env Health Perspectives* 1999; 107; Suppl 2:227-306.
- Boffetta P, Kogevinas M, Westerholm P, Saracci R. Exposure to occupational carcinogens and social class differences in cancer occurrence. *IARC Scientific Publ* 1997;138:331-41.
- Bollini P, Siem H. No real progress towards equity: health of migrants and ethnic minorities on the eve of the year 2000. *Soc Sci Med* 1995;41:819-28.
- Botman SL, Jack SS. Combining national health interview survey datasets: issues and approaches. *Stats in Med* 1995;14:669-77.
- Brackbill R, Frazier T, Shilling S. Smoking characteristics of US workers, 1978-1980. *Am J Indust Med*. 1988; 13: 5-41.
- Brackbill RM, Cameron LL, Behrens V. Prevalence of chronic diseases and impairments among US farmers, 1986-90. *Am J Epid* 1994;139:1055-65.
- Burnett CA, Crouse WE. Agreement of latest and longest occupation and industry as reported by 1980 National Health Interview Survey. *JOM* 1989;31(4):339-346.
- Caban A, Fleming LE, Lee D, Gomez-Marin O, LeBlanc W, Pitman T. Obesity in U.S. Workers: The National Health Interview Survey 1986 – 2002. *J APHA* in press
- Carpenter L, Roman E. Cancer and occupation in women: identifying associations using routinely collected national data. *Environmental Health Perspectives*. 1999; 107 Suppl 2:299-303.

- Cassano C. de Medeiros Frias LA, Valente JG. *Cadernos de Saude Publica*. 2000;16(## Suppl 1):53-64.
- Checkoway H, Pearce NE, Kriebel D. *Research Methods in Occupational Epidemiology*, Second Edition. New York: Oxford University Press, 2004.
- Chiazze L, Watkins DK, Fryar C. Historical cohort mortality study of a continuous filament fiberglass manufacturing plant. *JOEM* 1997;39:432-41.
- Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas* 1960; 20:37– 46.
- Cooper SP, Buffler PA, Lee ES, Cooper CJ. Health characteristics by longest held occupation and industry of employment: US 1980. *Am J Ind Med* 1993;24:25-39.
- Dell L, Teta MJ. Mortality among workers at a plastics manufacturing and research and development facility. *Am J Ind Med* 1995;28:325-37.
- Dong X, Platner JW. Occupational fatalities of Hispanic construction workers from 1992 to 2000. *American Journal of Industrial Medicine*. 2004;45(1):45-54.
- Drever F ed. *The Registrar General's Decennial Supplement for England & Wales; Occupational Health: Decennial Supplement*. London: Office of Population Censuses and Surveys Health and Safety Executive, Series DS #10, 1995.
- Edwards WS, Kurlantzick V, et al. Evaluation of National Health Interview Survey diagnostic reporting. *Vital Health Stat 2*. Feb 1994(120):1-116.
- Edwards WS, Winn DM, Collins JG. Evaluation of 2-week doctor visit reporting in the national health interview survey. *Vital Health Stat 2*. Aug 1996(122):1-46.
- Engholm G, England A. Morbidity and mortality patterns in Sweden. *Occ Med* 1995;10:261-8.
- Fleming LE, Bean JA, Rudolph M, Hamilton K. Mortality in Florida Pesticide Applicators. *Occup Env Med* 1999:14-8.
- Fleming LE, Gomez-Marin O, Zheng D, Ma F, Lee D. National Health Interview Survey (NHIS) Mortality among US Farmers and Pesticide Applicators. *Am J Ind Med*, 2003;43(2):227-233.
- Fleming LE, LeBlanc W, Caban A, Pitman T, Lee DJ, Gomez Marin O. *Interactive Monograph: Occupation, Disability and Self Experienced Health in the National Health: 1986-1994*. 2004. <http://www.UmiamiORG.com>
- Fosbroke DE, Kisner SM, Myers JR. Working lifetime risk of occupational fatal injury. *Am J Ind Med* 1997;31:459-67.
- Frumkin H, Pransky G. Special Populations in occupational health. *Occ Med* 1999;14:479-84.
- Gallagher RP, Threlfall WJ, Band PR, Spinelli JJ. *Occupational Mortality in British Columbia 1950-1984*. Vancouver, BC: Cancer Control Agency of British Columbia, 1989.
- Gregoire A. The mental health of farmers. *Occupational Medicine (Oxford)*. 2002;52(8):471-6.

Gold B, Kathren RL. Causes of death of a cohort of 260 plutonium workers. *Health Phys* 1998;75:236-40.

Gomez Marin O, Fleming LE, Caban A, LeBlanc W, Lee DJ, Pitman T. Longest Held Job in US Occupational Groups: The National Health Interview Survey (NHIS). *Journal Occupational and Environmental Health* 2005;47(1):79-90.

Gomez Marin O, Fleming LE, Lee DJ, LeBlanc W, Zheng D, Ma F, Jane D, Pitman T, Caban A. Acute and Chronic Disability among US Farmers and Pesticide Applicators: The National Health Interview Survey (NHIS). *Journal Agricultural Safety Health* 2004; 10(4): 285-293.

Guralnick L. Mortality by industry and cause of death among men 20-64 years of Age: US, 1950. Vol 53(4). Washington DC: NCHS, 1963b.

Guralnick L. Mortality by occupation and causes of death among men 20-64 years of Age: US, 1950. Vol 53(3). Washington DC: NCHS, 1963a.

Guralnick L. Mortality by occupation and industry among men 20-64 years of Age: US, 1950. Vol 53(2). Washington DC: NCHS, 1962.

Healthy People 2010 Statistical Notes #20: Age Adjustment Using the 2000 Projected U.S. Population. 10 pp. (PHS) 2001-1237. <http://www.cdc.gov/nchs/products/pubs/pubd/hp2k/statnt/20-11.htm>

Honda Y, Delzell E, Cole P. An updated study of mortality among workers at a petroleum manufacturing plant. *JOEM* 1995;37:194-200.

Hwang SA, Fitzgerald EF, Herzfeld PM, Stark A. Mortality among NY highway maintenance workers. *Int Arch Occ Env Health* 1995;67:225-35.

Janicak CA. Differences in relative risks for fatal occupational highway transportation accidents. *Journal of Safety Research*. 2003;34(5):539-45.

Janzen BL, Muhajarine N. Social role occupancy, gender, income adequacy, life stage, and health: a longitudinal study of employed Canadian men and women. *Soc Sci Med* 2003;57:1491-1503.

Kagamimori S, Matusbara I, Sokejima S, et al. The comparative study of occupational mortality between Japan and Great Britain. *Ind Health* 1998;36:252-7.

Kaminski R, Spirtas R. *Industrial Characteristics of Persons Reporting Morbidity during the Health Interview Surveys Conducted in 1969-1974*. Cincinnati, OH: NIOSH, 1980.

Khlat M, Sermet C, Le Pape A. Women's health in relation with their family and work roles, France in the early 1990s. *Soc Sci Med* 2000;50:1807-1825.

King TK, Borrelli B, Black C, Pinto BM, Marcus BH. Minority women and tobacco: implications for smoking cessation interventions. *Ann Behav Med* 1997;19:301-13.

Kircher T, Nelson J, Burdo H. The autopsy as a measure of accuracy of the death certificate. *NEJM* 1985;313:1263-1269.

Kirkhorn SR, Schenker MB. Current health effects of agricultural work: respiratory disease, cancer,

reproductive effects, musculoskeletal injuries, and pesticide-related illnesses. *Journal of Agricultural Safety & Health*. 2002;8(2):199-214.

Koskela RS. Mortality, morbidity, and health selection among metal workers. *Scand J Work Env Health* 1997;23 Suppl 2:1-80.

Kreiss K, Cox-Ganser J. Metalworking fluid-associated hypersensitivity pneumonitis: a workshop summary. *American Journal of Industrial Medicine*. 1997;32(4):423-32.

Kross BC, Burmeister LF, Ogilvie LK, Fuortes LJ, Fu CM. Proportionate mortality study of golf course superintendents. *Am J Ind Med* 1996;29:501-6.

Lam LT. Environmental factors associated with crash-related mortality and injury among taxi drivers in New South Wales, Australia. *Accident Analysis & Prevention*. 2004;36(5):905-8.

Lee DJ, LeBlanc W, Fleming LE, Gomez-Marin O, Pitman T. Trends in US Smoking Rates in Occupational Groups: The National Health Interview Survey 1987-1994. *JOEM* 2004;46:538-548.

Liao Y, Cooper RS, Cao G et al. Mortality patterns among adult Hispanics: findings from the NHIS, 1986-1990. *Am J Pub Health* 1998;88:227-32.

Loomis D, Richardson D. Race and the risk of fatal injury at work. *Am J Pub Health* 1998;88:40-4.

Ma F, Lee DJ, Fleming LE, Mustafa D. Race specific cancer mortality in US Firefighters: 1984-1993. *J Occ Env Med* 1998;40:1134-8.

Martins PJ, D'Almeida V, Vergani N, Perez AB, Tufik S. Increased plasma homocysteine levels in shift working bus drivers. *Occupational & Environmental Medicine*. 2003;60(9):662-6.

Massey JT, Morre TF, Parson VL, Tadros W. Design and Estimation for the NHIS 1985-94. Washington DC: NCHS, Vital & Health Statist Series 2#110 DHHS Pub PHS89-1384, 1989.

Matanoski GW, Kanchanaraks S, Lees PS, et al. Industry wide study of mortality of pulp and paper mill workers. *Am J Ind Med* 1998;33:327-40.

Melius J. Asphalt--a continuing challenge. *American Journal of Industrial Medicine*. 2003;43(3):235-6.

Milham S. Occupational Mortality in Washington State 1950-71. Cincinnati, OH: NIOSH, Dept of HEW #76-175-A, 1976.

Milham S. Occupational Mortality in Washington State 1950-79. Cincinnati, OH: NIOSH, Dept of HEW #83-116, 1983.

Milham S. Cancer among farmers: a meta-analysis. *Annals of Epidemiology*. 1999;9(1):71-2.

Monson RR. *Occupational Epidemiology*. Boca Raton: CRC Press, 1990, second edition.

Murray LR. Sick and tired of being sick and tired: scientific evidence, methods, and research implications for racial and ethnic disparities in occupational health. *AM J Pub Health* 2003;93:221-226.

Nathell L, Malmberg P, Lundback B, Nygren A. Impact of occupation on respiratory disease.

Scandinavian Journal of Work, Environment & Health. 2000;26(5):382-9.

National Center for Health Statistics (NCHS). Classified index of industries and occupations, 1989. Washington DC: US Dept of Health and Human Services. National Center for Health Statistics (NCHS), 1998a.

National Center for Health Statistics (NCHS). Documentation for the NHIS Multiple Cause of

Death Public Use Data File 1986-94 Survey Years; Dates of Death 1986-95. Washington, DC: US Dept of Health and Human Services, NCHS, 1998b.

National Center for Health Statistics (NCHS). Documentations for the NHIS Surveys 2001. Washington, DC: US Dept of Health and Human Services, NCHS, 2001.

National Center for Health Statistics (NCHS). Industry and occupation coding for death certificates, 1993. NCHS instruction manual, Part 19. Hyattsville, Maryland: Public Health Service, 1992.

Nelson DE, Emont SL, Brackbill RM, Cameron LL, Peddicord J, Fiore MC. Cigarette smoking prevalence by occupation in the US. JOEM 1994;36:516-25.

National Institute of Occupational Safety and Health (NIOSH). Fatal Injuries to Workers in the US, 1980-89. Cincinnati, OH: NIOSH #93-108, 1993.

National Institute of Occupational Safety and Health (NIOSH). Mortality by Occupation, Industry, and Cause of Death: 24 Reporting States (1984-88). Cincinnati, OH: NIOSH #1997-114, 1997.

O'Campo P, Eaton WW, Muntaner C. Labor market experience, work organization, gender inequalities, and health status: results from a prospective analysis of US employed women. Soc Sci Med 2004;58:585-594.

Parron T. Hernandez AF. Villanueva E. Increased risk of suicide with exposure to pesticides in an intensive agricultural area. A 12-year retrospective study. Forensic Science International. 1996;79(1):53-63.

Partanen T. Johansson M. Ahrens W. et al. Assessment of feasibility of workplace health promotion. Preventive Medicine. 2002;35(3):232-40.

Plani F. Bowley DM. Goosen J. Death and injury on duty--a study of South African police officers. South African Medical Journal. Suid-Afrikaanse Tydskrif Vir Geneeskunde. 2003;93(11):851-3.

Pollan M, Gustavsson P. High risk occupations for breast cancer in the Swedish female working population. Am J Pub Health 1999;89:875-81.

Research Triangle Institute (RTI). Survey Data Analysis (SUDAAN). RTI, 2001.

Reviere R, Schneider S, Woolbright K. Associations between disease and occupation: hypotheses generated from the National Mortality Followback Survey. Am J Ind Med 1995;27:195-205.

Robinson CF, Halperin WE, Alterman T et al. Mortality patterns among construction workers in the US. Occ Med 1995;10:269-83.

Rosengren A. Wilhelmsen L. Cancer incidence, mortality from cancer and survival in men of different

occupational classes. *European Journal of Epidemiology*. 2004;19(6):533-40.

Rosenstock L, Cullen M, et al eds. *Textbook of Clinical Occupational and Environmental Medicine*, 2nd Ed. Orlando: Saunders, 2005,

Savitz DA, Checkoway H, Loomis DP. Magnetic field exposure and neurodegenerative disease mortality among electric utility workers. *Epid* 1998;9:398-404.

Schade WJ, Swanson GM. Comparison of death certificate occupation and industry data with lifetime occupational histories obtained by interview: variations in the accuracy of death certificate entries. *AM J Ind Med* 1988;14:121-136.

Schouten EJ, Borgdorff MW. Increased mortality among Dutch development workers. *BMJ* 1995;311:1343-4.

Schumacher MC. Comparison of occupation and industry information from death certificates and interviews. *AmJPH* 1986;76(6): 635-637.

Simpson J. Roman E. Law G. Pannett B. Women's occupation and cancer: preliminary analysis of cancer registrations in England and Wales, 1971-1990. *American Journal of Industrial Medicine*. 1999;36(1):172-85.

Steenland K, Brown D. Mortality study of goldminers exposed to silica and non-asbestiform amphibole minerals: an update with 14 more years of followup. *Am J Ind Med* 1995;27:195-205.

Steenland K, Burnett C, Lalich N, Ward E, Hurrell J. Dying for work: the magnitude of US mortality from selected causes of death associated with occupation. *Am J Ind Med* 2003b;43:461-482.

Steenland K, Halperin W, Hu S, Walker JT. Deaths due to injuries among employed adults: the effects of socio-economic class. *Epidemiology* 2003a;14:74-79.

Stein AD, Mincheva V, Stoyanovsky V. Trends in cardiovascular disease risk factor prevalence among male transport workers: Bulgaria, 1986 to 1997. *American Journal of Public Health*. 2001;91(3):455-7.

Sterling T, Weinkam J. The confounding of occupation and smoking and its consequences. *Soc Sci Med* 1990;30:457-67.

Stout NA, Jenkins EL, Pizatella TJ. Occupational injury mortality rates in the US: Changes from 1980-89. *Am J Public Health* 1996;86:73-7.

Suruda A, Wallace D. Fatal work related injuries in the US chemical industry 1984-89. *Int Arch Occ Env Health* 1996;68:425-8.

Thornberry O. An experimental comparison of telephone and personal health surveys. *Vital Health Stat 2*. 1987;106:1-186.

Toscano G, Jack T. Occupational injury fatalities -- 1994. *Statistical Bulletin -- Metropolitan Insurance Company* 1996;77:12-22.

Tsai SP, Gilstrap EL, Cowles SR, Snyder PJ, Ross CE. Long term follow-up mortality study of petroleum refinery and chemical plant employees. *Am J Ind Med* 1996;29:75-87.

Wagener DK, Walstedt J, Jenkins L, Burnett C, Lalich N, Fingerhut M. Women: work and health. *Vital Health Stats* 1997;31:1-91.

Wagener DK, Winn DW. Injuries in working populations: black-white differences. *Am J Public Health* 1991;81:1408-14.

Zwerling C, Whitten PS, Davis CS, Sprince NL. Occupational injuries among workers with disabilities: the NHIS 1985-94. *JAMA* 1997;278:2163-6.

Zwerling C, Whitten PS, Davis CS, Sprince NL. Occupational injuries among older workers with visual, auditory, and other impairments. A validation study. *J Occ Env Med* 1998;40:720-3.

Zwerling C, Sprince NL, Wallace RB, Davis CS, Whitten PS, Geeringa SG. Risk Factors for Occupational Injuries among Older Workers: An Analysis of the Health and Retirement Study. *Am J Pub Health* 1996;86:1306-9.

Appendix

[Table 1](#) . 1986-1994 CDC Top 10 Causes of Death, ICD 9 Codes, and Deaths among Survey Respondents (Download [Table 1](#) in Excel Format)

[Table 2](#) . Occupational Groups, Number of Deaths, Sample Sizes, and Population Estimates by [13](#), [41](#) and [206 Occupations](#) (Download [Table 2](#) in Excel Format)

[Table 3](#) . Age Adjusted Overall and Cause Specific Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) (Download [Table 3](#) in Excel Format)

[Table 4](#) . Crude and Age Adjusted All Cause Mortality Rate/100,000 Person Years and SE among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 4](#) in Excel Format)

[Table 5](#) . Crude and Age Adjusted Cardiovascular Disease Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 5](#) in Excel Format)

[Table 6](#) . Crude and Age Adjusted Malignant Neoplasms Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 6](#) in Excel Format)

[Table 7](#) . Crude and Age Adjusted Cerebrovascular Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 7](#) in Excel Format)

[Table 8](#) . Crude and Age Adjusted Chronic Respiratory Disease Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 8](#) in Excel Format)

[Table 9](#) . Crude and Age Adjusted Unintentional Injury Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 9](#) in Excel Format)

[Table 10](#) . Crude and Age Adjusted Pneumonia and Influenza Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 10](#) in Excel Format)

[Table 11](#) . Crude and Age Adjusted Diabetes Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 11](#) in Excel Format)

[Table 12](#) . Crude and Age Adjusted Suicide Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 12](#) in Excel Format)

[Table 13](#) . Crude and Age Adjusted Liver Disease Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206 Occupations](#) by Socio-Demographic Subgroups (Download [Table 13](#) in Excel Format)

Table 14 . Crude and Age Adjusted HIV/AIDS Mortality Rate/100,000 Person Years (\pm SE) among [13](#), [41](#) and [206](#) Occupations by Socio-Demographic Subgroups (Download [Table 14](#) in Excel Format)